

quences. Most of these layers were deposited under quiet, low-energy, subaqueous conditions. In shallow-water sections, they appear to include both *in situ* silicified bacterial mats and detritus eroded from them. Carbonaceous matter in deeper-water deposits consists exclusively of fine-grained pelagic, hemipelagic, or current-deposited detritus.

Preserved early Archean stromatolites and carbonaceous matter appear to reflect communities of photosynthetic cyanobacteria inhabiting shallow, probably marine environments developed over the surfaces of low-relief, rapidly subsiding, simatic volcanic platforms. The overall environmental and tectonic conditions were those that probably prevailed at Earth's surface since the simatic crust and oceans formed sometime before 3,800 Ma. Recent studies also suggest that these early Archean sequences contain layers of debris formed by large-body impacts on early Earth (Lowe and Byerly, 1986, 1988; Lowe *et al.*, 1988). If so, then these early bacterial communities had developed strategies for coping with the disruptive effects of possibly globe-encircling high-temperature impact vapour clouds, dust blankets, and impact-generated tsunamis. It is probable that these early Archean biogenic materials represent organic communities that evolved long before the beginning of the preserved geologic record and were well adapted to the rigors of life on a young, volcanically active Earth during late bombardment. These conditions may have had parallels on Mars during its early evolution.

REFERENCES

- Byerly, G. R., Lowe, D. R., and Walsh, M. M. Stromatolites from the 3,300-3,500-Myr Swaziland Supergroup, Barberton Mountain Land, South Africa. *Nature* **319**:489-491, 1986.
- Lowe, D. R. Stromatolites 3,400-Myr old from the Archean of Western Australia. *Nature* **284**:441-443, 1980.
- Lowe, D. R. Early Archean organisms and paleoenvironments: 5th meeting of the International Soc. for the Study of the Origin of Life, Berkeley, Abstracts, p. 7, 1986.
- Lowe, D. R., and Byerly, G. R. Early Archean spherules of probable impact origin, South Africa and Western Australia. *Geology* **14**:83-86, 1986.
- Lowe, D. R., and Byerly, G. R. Identification and effects of large, early Archean, terrestrial meteorite impacts: a geological perspective on late accretion. 19th Lunar and Planetary Science Conference, Abstracts, 1988.
- Lowe, D. R., Asaro, F., and Byerly, G. R. Iridium anomalies associated with early Archean spherule layers, South Africa and Western Australia. 19th Lunar and Planetary Science Conference, Abstracts, 1988.
- Walter, M. R., Buick, R., and Dunlop, J. S. R. Stromatolites 3,400-3,500 Myr old from the North Pole area, Western Australia. *Nature* **284**:443-445, 1980.